



## Anti-VACHT antibody (STJ13102599)

## GENERAL INFORMATION

|               |                    |
|---------------|--------------------|
| Product Type  | Primary antibodies |
| Applications  | IHC/WB             |
| Host / Source | Goat               |
| Reactivity    | Human/Mouse/Rat    |

## PRODUCT PROPERTIES

|                     |                                                                                                                                                                                                                                                   |
|---------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Clonality           | Polyclonal                                                                                                                                                                                                                                        |
| Conjugation         | Unconjugated                                                                                                                                                                                                                                      |
| Purification        | IgG purified                                                                                                                                                                                                                                      |
| Dilution Range      | IHC, WB. A working concentration of 10-50 ug/ml is recommended. The optimal concentration should be determined by the end user. Not yet tested in other applications.                                                                             |
| Formulation         | Lyophilised                                                                                                                                                                                                                                       |
| Isotype             | IgG                                                                                                                                                                                                                                               |
| Storage Instruction | Maintain the lyophilised/reconstituted antibodies frozen at -20°C for long term storage and refrigerated at 2-8°C for a shorter term. When reconstituting, Glycerol (1:1) may be added for an additional stability. Avoid freeze and thaw cycles. |

## TARGET INFORMATION

|             |                     |
|-------------|---------------------|
| Specificity | Specific for VACHT. |
|-------------|---------------------|

## ADDITIONAL INFORMATION

|      |                                                                                                                         |
|------|-------------------------------------------------------------------------------------------------------------------------|
| Note | STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS. |
|------|-------------------------------------------------------------------------------------------------------------------------|