

Anti-MDM2 antibody [ZR258] (STJ180614)

GENERAL INFORMATION

| | |
|---------------|--------------------|
| Product Type | Primary antibodies |
| Applications | IHC-P |
| Host / Source | Rabbit |
| Reactivity | Human |

PRODUCT PROPERTIES


| | |
|---------------------|--|
| Clonality | Monoclonal |
| Conjugation | Unconjugated |
| Purification | Affinity purified |
| Dilution Range | 1:100-200 |
| Formulation | Tris-HCl buffer containing stabilizing protein (BSA) and <0.1% ProClin |
| Isotype | IgG |
| Storage Instruction | Store at 2-8°C for up to 24 months. Predilute: Ready to use, no reconstitution necessary. Concentrate: Use dilution range and appropriate lab-standardized diluent. Stability after dilution: 7 days at 24°C, 3 months at 2-8°C, 6months at -20°C. |

TARGET INFORMATION

| | |
|-------------|-------------------------------|
| Gene ID | 4193 |
| Gene Symbol | MDM2 |
| UniProt ID | MDM2_HUMAN |
| Specificity | Positive Control: Liposarcoma |

ADDITIONAL INFORMATION

Note STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.



Anti-MDM2 antibody IHC human atypical lipomatous tumor STJ180614