Anti-CASR antibody (180-536) (STJ22909)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/IHC-P/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES

Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:1000 IHC-P:1:50-1:200 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.09% Sodium Azide, 50% Glycerol, pH7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 121kDa Observed Mw: 130-260kDa
Storage Instruction	Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID	846
Gene Symbol	CASR
UniProt ID	CASR_HUMAN
Immunogen Region	180-536
Immunogen Sequence	FKSFLRTIPNDEHQATAMAD IIEYFRWNWVGTIAADDDYG RPGIEKFREEAEERDICIDF SELISQYSDEEEIQHVVEVI QNSTAKVIVVFSSGPDLEPL IKEIVRRNITGKIWLASEAW ASSSLIAMPQYFHVVGGTIG FALKAGQIPGFREFLKKVHP RKSVHNGFAKEFWEETFNCH LQEGAKGPLPVDTFLRGHEE SGDRFSNSSTAFRPLCTGDE NISSVETPYIDYTHLRISY
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 180-536 of human CASR (NP_000379.3).

ADDITIONAL INFORMATION

Note	STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.



Immunohistochemistry analysis of paraffin-embedded rat kidney using CASR antibody (STJ22909) at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6. 0 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse kidney using CASR antibody (STJ22909) at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6. 0 before commencing with immunohistochemistry staining protocol.



Western blot analysis of extracts of various cell lines, using CASR antibody (STJ22909) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 180s.