



Anti-CHRM1 antibody (251-350) (STJ23130)


GENERAL INFORMATION		
	Product Type	Primary antibodies
	Applications	WB/IHC-P/IF/ICC/ELISA
	Host / Source	Rabbit
	Reactivity	Human/Mouse/Rat
PRODUCT PROPERTIES		
	Clonality	Polyclonal
	Concentration	Lot specific
	Conjugation	Unconjugated
	Purification	Affinity purification
	Dilution Range	WB:1:500-1:1000 IHC-P:1:50-1:200 IF/ICC:1:50-1:200 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
	Formulation	PBS with 0.09% Sodium Azide, 50% Glycerol, pH7.3.
	Isotype	IgG
	Molecular Weight	Protein Mw: 51kDa Observed Mw: 51kDa
	Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.
TARGET INFORMATION		
	Gene ID	1128
	Gene Symbol	CHRM1
	UniProt ID	ACM1_HUMAN
	Immunogen Region	251-350
	Immunogen Sequence	SPETPPGRCCRCRAPRLQ AYSWKEEEEDEGSMESLTS SEGEEPGSEVVIKMPMVDPE AQAPTQPPRSSPNTVKRPT KKGDRDAGKGQKPRGKEQLA
	Specificity	A synthetic peptide corresponding to a sequence within amino acids 251-350 of human CHRM1 (NP_000729.2).
ADDITIONAL INFORMATION		
	Note	STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.




Immunofluorescence analysis of SH-SY5Y cells using CHRM1 Rabbit polyclonal antibody (STJ23130) at dilution of 1:200 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of Neuro-2a cells using CHRM1 Rabbit polyclonal antibody (STJ23130) at dilution of 1:200 (40x lens). Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of paraffin-embedded mouse brain using CHRM1 Rabbit polyclonal antibody (STJ23130) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with immunohistochemistry staining protocol.



Western blot analysis of extracts of various cell lines, using CHRM1 antibody (STJ23130) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 μ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 30s.