

Anti-ERG antibody (1-280) (STJ23567)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/ELISA
Host / Source	Rabbit
Reactivity	Human

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 54kDa Observed Mw: 60kDa
Storage Instruction	Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID	2078
Gene Symbol	ERG
UniProt ID	ERG_HUMAN
Immunogen Region	1-280
Immunogen Sequence	MASTIKEALSVMSEDQSLFE CAYGTPHLAKTEMTASSSSD YGQTSKMSPRVPQQDWLSQP PARVTIKMECNPSQVNGSRN SPDECSVAKGGKMGSPDTV GMNYGSYMEEKHMPPPNMTT NERRVIVPADPTLWSTDHVR QWLEWAVKEYGLPDVNILLF QNIDGKELCKMTKDDFQRLT PSYNADILLSHLHYLRETP LPHLTSDDDVKALQNSPRLMH ARNTGGAAFIFPNTSVYPE
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 1-280 of human ERG (NP_891548.1).

ADDITIONAL INFORMATION

Note	STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.
------	---



Western blot analysis of extracts of various cell lines, using ERG antibody (STJ23567) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST.