

Anti-MYH9 antibody (1711-1960) (STJ24664)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/IHC-P/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:1000-1:2000 IHC-P:1:50-1:100 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 227kDa Observed Mw: 250kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	4627
Gene Symbol	MYH9
UniProt ID	MYH9_HUMAN
Immunogen Region	1711-1960
Immunogen Sequence	ANSSGKGGALALEEKRRLEAR IAQLEEELEEEQGNTELIND RLKKNALQIDQINTDLNLER SHAQKNENARQQLERQNKEL KVKLQEMEGTVKSKYKASIT ALEAKIAQLEEQLDNETKER QAACKQVRRTEKKLKDVLQ VDDERRNAEQYKDQADKAST RLKQLKRQLEEEAEEEAQRAN ASRRKLQRELEDATETADAM NREVSSLKNKLRRGDLPFVV PRRMARKGAGDGSDEEVDG
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 1711-1960 of human MYH9 (NP_002464.1).

ADDITIONAL INFORMATION


Note **STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.**



Immunohistochemistry analysis of paraffin-embedded human stomach using MYH9 antibody (STJ24664) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of paraffin-embedded human esophagus using MYH9 antibody (STJ24664) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Western blot analysis of extracts of various cell lines, using MYH9 antibody (STJ24664) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 15s.