

Anti-MYOC antibody (245-504) (STJ24669)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 57kDa Observed Mw: 65kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID	4653
Gene Symbol	MYOC
UniProt ID	MYOC_HUMAN
Immunogen Region	245-504
Immunogen Sequence	CGELVWVGEPLTLRTAETIT GKYGVWMRDPKPTYPTQET TWRIDTVGTDVVRQVFEYDLI SQFMQGYPSKVHILPRPLES TGAVVYSGSLYFQGAESRTV IRYELNTETVKAEKEIPGAG YHGQFPYSWGGYTDIDLAVD EAGLWVIYSTDEAKGAVLS KLNPNLELEQTWETNIRKQ SVANAFICGTLTYVSSYTS ADATVNFAYDTGTGISKTLT IPFKNRYKYSSMIDYNPLE
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 245-504 of human MYOC (NP_000252.1).

ADDITIONAL INFORMATION

Note **STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.**



Western blot analysis of extracts of various cell lines, using MYOC antibody (STJ24669) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 60s.