Anti-STIP1 antibody (1-300) (STJ25731)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/IHC-P/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES

Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 IHC-P:1:50-1:200 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 63kDa Observed Mw: 63kDa
Storage Instruction	Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID	10963
Gene Symbol	STIP1
UniProt ID	STIP1_HUMAN
Immunogen Region	1-300
Immunogen Sequence	MEQVNELKEKGNKALSVGNI DDALQCYSEAIKLDPHNHVL YSNRSAAYAKKGDYQKAYED GCKTVDLKPDWGKGYSRKAA ALEFLNRFEEAKRTYEEGLK HEANNPQLKEGLQNMEARLA ERKFMNPFNMPNLYQKLESD PRTRTLLSDPTYRELIEQLR NKPSDLGTKLQDPRIMTTLS VLLGVDLGSMDEEEEIATPP PPPPPKKETKPEPMEEDLPE NKKQALKEKELGNDAYKKK
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 1-300 of human STIP1 (NP_006810.1).

ADDITIONAL INFORMATION

Note	STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.



Immunohistochemistry analysis of paraffin-embedded rat testis using STIP1 Rabbit polyclonal antibody (STJ25731) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9. 0 before commencing with immunohistochemistry staining protocol.



Western blot analysis of extracts of various cell lines, using STIP1 antibody (STJ25731) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST.