

Anti-TLR2 antibody (636-784) (STJ25865)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/IF/ICC/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:1000 IF/CC:1:50-1:200 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 90kDa Observed Mw: 105kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	7097
Gene Symbol	TLR2
UniProt ID	TLR2_HUMAN
Immunogen Region	636-784
Immunogen Sequence	SRNICYDAFVSYSERDAYWV ENLMVQELENFNPPFKLCLH KRDFIPGKWIIDNIDSIEK SHKTVFVLSSENFVKSEWCKY ELDFSHFRLFDENNDAAILI LLEPIEKKAIPQRFCCLRKI MNTKTYLEWPMDEAQRGEFW VNLRAAIKS
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 636-784 of human TLR2 (NP_003255.2).

ADDITIONAL INFORMATION


Note **STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.**



Immunofluorescence analysis of U937 cells using TLR2 Rabbit polyclonal antibody (STJ25865) at dilution of 1:200 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of Raw264 7 cells using TLR2 Rabbit polyclonal antibody (STJ25865) at dilution of 1:200 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Western blot analysis of various lysates using TLR2 Rabbit polyclonal antibody (STJ25865) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 1s.