

Anti-HMG20B antibody (1-317) (STJ26659)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:1000 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 36kDa Observed Mw: 36kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID	10362
Gene Symbol	HMG20B
UniProt ID	HM20B_HUMAN
Immunogen Region	1-317
Immunogen Sequence	MSHGPKQPGAAAAPAGGKAP GQHGGFVVTVKQEREGEPRA GEKGSHEEEPVKKRGWPKGK KRKKILPNGPKAPVTGYVRF LNERREQIRTRHPDLFPPEI TKMLGAEWSKLQPTKQRYL DEAEREKQQYMKELRAYQQS EAYKMCTEKIQEKKIKKEDS SSGLMNTLLNGHKGDCDGF STFDVPIFTEEFLDQNKARE AELRRLRKMNVAFEEQNAVL QRHTQSMSSARERLEQELA
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 1-317 of human HMG20B (NP_006330.2).

ADDITIONAL INFORMATION

Note	STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.
------	---



Western blot analysis of extracts of various cell lines, using HMG20B antibody (STJ26659) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% non-fat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.