

Anti-MCM5 antibody (1-230) (STJ27502)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/IF/ICC/IP/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:1000-1:3000 IF/ICC:1:50-1:200 IP:0.5 Mu g-4 Mu g antibody for 200 Mu g-400 Mu g extracts of whole cells ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirement
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 82kDa Observed Mw: 100kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	4174
Gene Symbol	MCM5
UniProt ID	MCM5_HUMAN
Immunogen Region	1-230
Immunogen Sequence	MSGFDDPGIFYSDSFGGDAQ ADEGQARKSQLQRRFKEFLR QYRVGTDRTGFTFKYRDELK RHYNLGEYWIEVEMEDLASF DEDLADYLYKQPAEHLQLE EAAKEVADEVTRPRPSGEEV LQDIQVMLKSDASPSSIRSL KSDMMSHLVKIPGIIIAASA VRAKATRISIQCRSCRNTLT NIAMRPGLEGYALPRKCNTD QAGRPKCPLDPYFIMPDCKC CVDFTLKLQ
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 1-230 of human MCM5 (NP_006730.2).

ADDITIONAL INFORMATION


Note	STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.
------	---



Immunoprecipitation analysis of 300 Mu g extracts of HeLa cells using 3 Mu g MCM5 antibody (STJ27502). Western blot was performed from the immunoprecipitate using MCM5 antibody (STJ27502) at a dilution of 1:1000.



Immunofluorescence analysis of U-2 OS cells using MCM5 Rabbit polyclonal antibody (STJ27502) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Western blot analysis of various lysates using MCM5 Rabbit polyclonal antibody (STJ27502) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 30s.