

Anti-CLCN2 antibody (550-770) (STJ27873)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 99kDa Observed Mw: 90-98kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID	1181
Gene Symbol	CLCN2
UniProt ID	CLCN2_HUMAN
Immunogen Region	550-770
Immunogen Sequence	PSLYDSIIRIKKLPYLPELG WGRHQQYRVRVEDIMVRDVP HVALSCTFRDLRLALHRTKG RMLALVESPEMILLGSIER SQVVALLGAQLSPARRRQHM QERRATQTSPLSDQEGPPTP EASVCFQVNTEDSAFPAARG ETHKPLKPALKRGPSVTRNL GESPTGSAESAGIALRSLFC GSPPEAAASEKLESCERKL KVRISLASDADLEGEMSPE E
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 550-770 of human CLCN2 (NP_001164560.1).

ADDITIONAL INFORMATION

Note	STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.
------	---



Western blot analysis of various lysates using CLCN2 Rabbit pAb (STJ27873) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.