

Anti-CLCN1 antibody (779-988) (STJ28306)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/IHC-P/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:1000 IHC-P:1:50-1:200 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.05% Proclin300, 50% Glycerol, pH7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 109kDa Observed Mw: 109kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	1180
Gene Symbol	CLCN1
UniProt ID	CLCN1_HUMAN
Immunogen Region	779-988
Immunogen Sequence	ARPTKKKTTQDSTDLDVNMS PEEIEAWEEQQLSQPVCFDS CCIDQSPFQLEQTTLHKTH TLFSLGLHLAYVTSMGKLR GVLALEELQKAIEGHTKSGV QLRPPLASFRNTTSTRKSTG APPSSAENWNLPEDRPGATG TGDVIAASPETVPVSPSPSEP PLSLAPGKVEGELEELVE SPGLEEELADILQGPSLRST DEEDEDELIL
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 779-988 of human CLCN1 (NP_000074.3).

ADDITIONAL INFORMATION


Note	STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.
------	---



Immunohistochemistry analysis of paraffin-embedded rat skeletal muscle using CLCN1 Rabbit polyclonal antibody (STJ28306) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse skeletal muscle using CLCN1 Rabbit polyclonal antibody (STJ28306) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with immunohistochemistry staining protocol.



Western blot analysis of extracts of Mouse skeletal muscle, using CLCN1 antibody (STJ28306) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 180s.