

Anti-TRIM32 antibody (1-280) (STJ29159)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:1000 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 72kDa Observed Mw: 75kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	22954
Gene Symbol	TRIM32
UniProt ID	TRI32_HUMAN
Immunogen Region	1-280
Immunogen Sequence	MAAAASHLNLDALREVLEC PICMESFTEEQLRPKLLHCG HTICRQCLEKLASSINGVR CPFCSKITRITSLTQLTDNL TVLKIIDTAGLSEAVGLLMC RSCGRRRLPRQFCRSCGLVLC EPCREADHQPPGHCTLPVKE AAEERRRDFGEKLTRLRLM GELQRRKAALGVSKDLQAR YKAVLQEYGHERRVQDELA RSRKFFTGS�AEVEKSNSQV VEEQSYLLNIAEVQAVSRC
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 1-280 of human TRIM32 (NP_036342.2).

ADDITIONAL INFORMATION

Note **STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.**



Western blot analysis of extracts of HeLa cells, using TRIM32 antibody (STJ29159) at 1:500 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 30s.



Western blot analysis of extracts of various cell lines, using TRIM32 antibody (STJ29159) at 1:500 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 10s.