

Anti-BCAS3 antibody (674-913) (STJ29413)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/IHC-P/IF/ICC/IP/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 IHC-P:1:50-1:200 IF/CC:1:50-1:200 IP:0.5 Mu g-4 Mu g antibody for 200 Mu g-400 Mu g extracts of whole cells ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specif
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 101kDa Observed Mw: 115kDa
Storage Instruction	Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	54828
Gene Symbol	BCAS3
UniProt ID	BCAS3_HUMAN
Immunogen Region	674-913
Immunogen Sequence	LAGLVPPGSPGPITRHGSYD SLASDHSQGEDEEWLSQVEI VTHTGPHRRLWMGPQFQFKT IHPSGQTTVISSSSVLQSH GPSDTPQPLDFDITDDLDLN SLRIQPVRSDPVSMPGSSRP VSDRRGVSTVIDAASGTFDR SVTLLEVCGSWPEGFGLRHM SSMEHTEEGLRERLADAMAE SPSRDVGSGTELRQREGSIE TLSNSSGSTSGSIPRNFDFGY RSPLEPTNESQLSLFPTGF
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 674-913 of human BCAS3 (NP_060149.3).

ADDITIONAL INFORMATION


Note STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.




Immunoprecipitation analysis of 200 Mu g extracts of HeLa cells, using 3 Mu g BCAS3 antibody (STJ29413). Western blot was performed from the immunoprecipitate using BCAS3 antibody (STJ29413) at a dilution of 1:1000.




Immunofluorescence analysis of A-549 cells using BCAS3 Rabbit polyclonal antibody (STJ29413). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunohistochemistry analysis of BCAS3 in paraffin-embedded mouse kidney using BCAS3 Rabbit polyclonal antibody (STJ29413) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7. 2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of BCAS3 in paraffin-embedded human stomach using BCAS3 Rabbit polyclonal antibody (STJ29413) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7. 2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of BCAS3 in paraffin-embedded human kidney cancer using BCAS3 Rabbit polyclonal antibody (STJ29413) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7. 2 before commencing with immunohistochemistry staining protocol.



Western blot analysis of various lysates using BCAS3 Rabbit polyclonal antibody (STJ29413) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.