Anti-NAA40 antibody (1-237) (STJ29522)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/ELISA
Host / Source	Rabbit
Reactivity	Human

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:1000 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 27kDa Observed Mw: 27kDa
Storage Instruction	Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID	79829
Gene Symbol	NAA40
UniProt ID	NAA40_HUMAN
Immunogen Region	1-237
Immunogen Sequence	MGRKSSKAKEKKQKRLEERA AMDAVCAKVDAANRLGDPLE AFPVFKKYDRNGLNVSIECK RVSGLEPATVDWAFDLTKTN MQTMYEQSEWGWKDREKREE MTDDRAWYLIAWENSSVPVA FSHFRFDVECGDEVLYCYEV QLESKVRRKGLGKFLIQILQ LMANSTQMKKVMLTVFKHNH GAYQFFREALQFEIDDSSPS MSGCCGEDCSYEILSRRTKF GDSHHSHAGGHCGGCCH
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 1-237 of human NAA40 (NP_079047.2).

ADDITIONAL INFORMATION

Note	STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.



Western blot analysis of lysates from 293T, using NAA40 rabbit polyclonal antibody (STJ29522) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% non-fat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 180s.