

## Anti-NAA50 antibody (1-169) (STJ29524)

### GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/IF/ICC/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

### PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 IF/CC:1:50-1:200 ELISA:Recommended starting concentration is 1 $\mu$ g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 19kDa Observed Mw: 19kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

### TARGET INFORMATION


Gene ID	<a href="#">80218</a>
Gene Symbol	<a href="#">NAA50</a>
UniProt ID	<a href="#">NAA50_HUMAN</a>
Immunogen Region	1-169
Immunogen Sequence	MKGSRIELGDVTPHNIKQLK RLNQVIFPVSYNDKFYKDVL EVGELAKLAYFNDAVAVGVC CRVDHSQNQKRLYIMTLGCL APYRRLGIGTKMLNHVLNIC EKDGTDFDNIYLHVQISNESA IDFYRKFGFEIETKKNYYK RIEPADAHVLRQKMLKVPVSGQ NADVQKTDN
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 1-169 of human NAA50 (NP_079422.1).

### ADDITIONAL INFORMATION

Note **STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.**



Immunofluorescence analysis of A549 cells using NAA50 antibody (STJ29524). Blue: DAPI for nuclear staining.



Western blot analysis of extracts of various cell lines, using NAA50 antibody (STJ29524) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25  $\mu$ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 90s.