Anti-POC1A antibody (138-407) (STJ29601)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 45kDa Observed Mw: 45kDa
Storage Instruction	Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID	25886
Gene Symbol	POC1A
UniProt ID	POC1A_HUMAN
Immunogen Region	138-407
Immunogen Sequence	LFSLSQHINWVRCAKFSPDG RLIVSASDDKTVKLWDKSSR ECVHSYCEHGGFVTYVDFHP SGTCIAAAGMDNTVKVWDVR THRLLQHYQLHSAAVNGLSF HPSGNYLITASSDSTLKILD LMEGRLLYTLHGHQGPATTV AFSRTGEYFASGGSDEQVMV WKSNFDIVDHGEVTKVPRPP ATLASSMGNLPEVDFPVPPG RGRSVESVQSQPQEPVSVPQ TLTSTLEHIVGQLDVLTQT
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 138-407 of human POC1A (NP_056241.3).

ADDITIONAL INFORMATION

Note	STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.



Western blot analysis of extracts of various cell lines, using POC1A antibody (STJ29601) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.