

Anti-SLC6A2 antibody (30-235) (STJ29850)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:1000 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.05% Proclin300, 50% Glycerol, pH7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 69kDa Observed Mw: 80kDa
Storage Instruction	Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID	6530
Gene Symbol	SLC6A2
UniProt ID	SC6A2_HUMAN
Immunogen Region	30-235
Immunogen Sequence	ELIPDAPLSSAAYSIRSIGE RPVLKAPVPKRQKCDHWTPC PSDTYAYRLLSGGGRSKYAK ICFEDNLLMGEQLGNVARGI NIAIVNYVTGNVTATRCFDM YEGDNSGPMTKFIQSAAPKS LLFMVTYDDGSTRLNNDAKN AIEALGSKEIRNMKFRSSWV FIAAKGLELPSEIQREKINH SDAKNNRYSGWPAEIQIEGC IPKERS
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 30-235 of human SLC6A2 (NP_478066.3).

ADDITIONAL INFORMATION

Note **STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.**



Western blot analysis of extracts of various cell lines, using SLC6A2 antibody (STJ29850) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 1s.