



Anti-SELP antibody (30-200) (STJ29851)


GENERAL INFORMATION		
	Product Type	Primary antibodies
	Applications	WB/IHC-P/ELISA
	Host / Source	Rabbit
	Reactivity	Human/Rat
PRODUCT PROPERTIES		
	Clonality	Polyclonal
	Concentration	Lot specific
	Conjugation	Unconjugated
	Purification	Affinity purification
	Dilution Range	WB:1:500-1:2000 IHC-P:1:50-1:200 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
	Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
	Isotype	IgG
	Molecular Weight	Protein Mw: 91kDa Observed Mw: 115kDa
	Storage Instruction	Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.
TARGET INFORMATION		
	Gene ID	6403
	Gene Symbol	SELP
	UniProt ID	LYAM3_HUMAN
	Immunogen Region	30-200
	Immunogen Sequence	ISELTNQKEVAAWTYHYSTK AYSWNISRKYQNRYTDLVA IQNKNEIDYLNKVLPPYSSY YWIGIRKNNKTWTWVGTKKA LTNEAENWADNEPNNKRNE DCVEIYIKSPSAPGKWDEH CLKKKHALCYTASCQDMSCS KQGECLTIGNYTCSCYPGF YGPECEYVREC
	Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 30-200 of human CD62P/P-selectin (NP_002996.2).
ADDITIONAL INFORMATION		
	Note	STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.




Immunohistochemistry analysis of paraffin-embedded human stomach using CD62P/P-selectin antibody (STJ29851) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of paraffin-embedded human esophageal cancer using CD62P/P-selectin antibody (STJ29851) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of paraffin-embedded human prostate using CD62P/P-selectin antibody (STJ29851) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Western blot analysis of extracts of various cell lines, using CD62P/P-selectin antibody (STJ29851) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 10s.