

## Anti-VPS28 antibody (C-Term) (STJ70508)

### GENERAL INFORMATION

Product Type	Primary antibodies
Applications	Pep-ELISA/WB/IHC
Host / Source	Goat
Reactivity	Human/Mouse/Rat/Dog/Pig

### PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	0.5 mg/mL
Conjugation	Unconjugated
Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Dilution Range	Peptide ELISA: antibody detection limit dilution 1:1000. WB: Approx 24kDa band observed in H460, Human Testis and Brain lysates (predicted MW of 25kDa according to NP_057292.1). Recommended for use at 1-3µg/ml. IHC: In paraffin embedded Human
Formulation	0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. NA
Isotype	IgG
Storage Instruction	Store at -20°C on receipt and minimise freeze-thaw cycles.

### TARGET INFORMATION


Gene ID	<a href="#">51160</a>
Gene Symbol	<a href="#">VPS28</a>
UniProt ID	<a href="#">VPS28_HUMAN</a>
Immunogen Region	C-Term
Immunogen Sequence	ESAYNAFNRLHA
Specificity	This antibody is expected to recognize isoform 1 (NP_057292.1) only.

### ADDITIONAL INFORMATION

Note **STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.**



STJ70508 staining (1µg/ml) of Human Testis lysate (RIPA buffer, 35µg total protein per lane). Primary incubated for 1 hour. Detected by chemiluminescence.



STJ70508 (10µg/ml) staining of paraffin embedded Human Testis. Microwaved antigen retrieval with Tris/EDTA buffer pH9, HRP-staining.