

## Anti-SNX26/TCGAP antibody (N-Term) (STJ70984)

### GENERAL INFORMATION

|               |                    |
|---------------|--------------------|
| Product Type  | Primary antibodies |
| Applications  | Pep-ELISA/IHC      |
| Host / Source | Goat               |
| Reactivity    | Human/Mouse        |

### PRODUCT PROPERTIES


|                     |   |
|---------------------|---|
| Clonality           | Polyclonal  |
| Concentration       | 0.5 mg/mL   |
| Conjugation         | Unconjugated  |
| Purification        | Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.   |
| Dilution Range      | Peptide ELISA: antibody detection limit dilution 1:32000.<br>WB: Preliminary experiments gave an approx 35kDa band in human brain (hippocampus) lysates after 1µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature. |
| Formulation         | 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. NA  |
| Isotype             | IgG   |
| Storage Instruction | Store at -20°C on receipt and minimise freeze-thaw cycles.  |

### TARGET INFORMATION

|                    |                             |
|--------------------|-----------------------------|
| Gene ID            | <a href="#">115703</a>      |
| Gene Symbol        | <a href="#">ARHGAP33</a>    |
| UniProt ID         | <a href="#">RHG33_HUMAN</a> |
| Immunogen Region   | N-Term                      |
| Immunogen Sequence | ARSTDSLDPGE                 |

### ADDITIONAL INFORMATION

Note STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.



STJ70984 1, 25µg/ml) staining of paraffin embedded human prostate. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.