

## Anti-FTL antibody (C-Term) (STJ71798)

### GENERAL INFORMATION

Product Type	Primary antibodies
Applications	Pep-ELISA/WB/IHC
Host / Source	Goat
Reactivity	Human/Mouse/Rat/Dog/Cow

### PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	0.5 mg/mL
Conjugation	Unconjugated
Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Dilution Range	Peptide ELISA: antibody detection limit dilution 1:32000. WB: Approx. 20kDa band observed in Human Brain (Cerebellum) , Liver, Placenta and 22kDa in Mouse and Rat Brain lysates (calculated MW of 20.0kDa according to NP_000137.2). Recommended conc
Formulation	0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. NA
Isotype	IgG
Storage Instruction	Store at -20°C on receipt and minimise freeze-thaw cycles.

### TARGET INFORMATION


Gene ID	<a href="#">2512</a>
Gene Symbol	<a href="#">FTL</a>
UniProt ID	<a href="#">FRIL_HUMAN</a>
Immunogen Region	C-Term
Immunogen Sequence	GEYLFERLTLKHD

### ADDITIONAL INFORMATION


Note **STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.**



STJ71798 (0.3 µg/ml) staining of Human Brain (Cerebellum) (A) , Liver (B) , Human Placenta (C) NA NA NA 22kDa in Mouse Brain (D) , Rat Brain lysate (35 µg protein in RIPA buffer). Detected by chemiluminescence.



STJ71798 (3.8 µg/ml) staining of paraffin embedded Human Brain Cortex. Steamed antigen retrieval with citrate buffer pH 6, HRP-staining. This data is from a previous batch, not on sale.



STJ71798 (1.5 µg/ml) as the reporter with EB002025 as the capture rabbit antibody (5 µg/ml). This data is from a previous batch, not on sale.