

Anti-PARK7/DJ-1, biotinylated antibody (C-Term) {Biotin} (STJ73276)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	Pep-ELISA/WB
Host / Source	Goat
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	0.5 mg/mL
Conjugation	Biotin
Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Dilution Range	Peptide ELISA: antibody detection limit dilution 1:128000. WB: Approx 24kDa band observed in lysates of cell line Jurkat (calculated MW of 19.9kDa according to NP_009193.2). See non-biotinylated parental product's datasheet for further QC data. R
Formulation	0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. NA
Isotype	IgG
Storage Instruction	Store at -20°C on receipt and minimise freeze-thaw cycles.

TARGET INFORMATION

Gene ID	11315
Gene Symbol	PARK7
UniProt ID	PARK7_HUMAN
Immunogen Region	C-Term
Immunogen Sequence	AAQVKAPLVLKD.
Specificity	Reported variants represent identical protein: NP_009193.2, NP_001116849.1

ADDITIONAL INFORMATION

Note STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.



Biotinylated STJ70331 (0.2 µg/ml) staining of Jurkat lysate (35 µg protein in RIPA buffer), exactly mirroring its parental non-biotinylated product. Primary incubation was 1 hour. Detected by chemiluminescence, using streptavidin-HRP and using NAP blocker as a substitute for skimmed milk.