

## Anti-SPIC antibody (131-180 aa) (STJ95749)

### GENERAL INFORMATION

|               |                    |
|---------------|--------------------|
| Product Type  | Primary antibodies |
| Applications  | WB/ELISA/IHC       |
| Host / Source | Rabbit             |
| Reactivity    | Human/Rat/Mouse    |

### PRODUCT PROPERTIES


|                     |   |
|---------------------|---|
| Clonality           | Polyclonal  |
| Concentration       | 1 mg/mL   |
| Conjugation         | Unconjugated  |
| Purification        | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution Range      | WB 1:500-2000<br>IHC-P 1:50-300<br>ELISA 2000-20000   |
| Formulation         | Liquid in PBS containing 50% Glycerol, 0.5% BSA and 0.02% Sodium Azide.   |
| Isotype             | IgG   |
| Molecular Weight    | Observed: 29kD  |
| Storage Instruction | Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.                        |

### TARGET INFORMATION


|                  |   |
|------------------|---|
| Gene ID          | <a href="#">121599</a>  |
| Gene Symbol      | <a href="#">SPIC</a>  |
| UniProt ID       | <a href="#">SPIC_HUMAN</a>  |
| Immunogen Region | 131-180 aa  |
| Specificity      | Spi-C Polyclonal Antibody detects endogenous levels of Spi-C protein. |

### ADDITIONAL INFORMATION


Note **STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.**




Western blot analysis of various cells using Spi-C polyclonal antibody, cells nucleus extracted by Minute™ Cytoplasmic STJ95749



Western blot analysis of lysates from HepG2 and HeLa cells, using SPIC Antibody. The lane on the right was blocked with STJ95749



Western blot analysis of the lysates from Jurkat cells using SPIC antibody. STJ95749



Immunohistochemical analysis of paraffin-embedded human oophoroma. 1, Antibody was diluted at 1:200 (4°C overnight). 2, STJ95749