

Anti-GST-Tag antibody [1B10] (STJ96909)

GENERAL INFORMATION

| | |
|---------------|---------------------|
| Product Type | Primary antibodies |
| Applications | WB/ELISA/IP |
| Host / Source | Mouse |
| Reactivity | Species independent |

PRODUCT PROPERTIES


| | |
|---------------------|--|
| Clonality | Monoclonal |
| Conjugation | Unconjugated |
| Purification | The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen. |
| Dilution Range | WB 1:5000 IP 1:200-500 ELISA 1:5000-20000 |
| Formulation | Liquid in PBS pH7.4, 0.5% BSA, 0.02% Sodium Azide and 50% Glycerol. |
| Isotype | IgG1 |
| Storage Instruction | Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles. |

TARGET INFORMATION

| | |
|-------------|--|
| Specificity | The antibody detects C-terminal, internal, and N-terminal GST fusion proteins. |
|-------------|--|

ADDITIONAL INFORMATION

Note STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.



Wang, YY., Zhao, B., Wu, MM. et al. Overexpression of neuregulin 1 in GABAergic interneurons results in reversible cortical disinhibition. Nat Commun 12, 278 (2021).
0.5 μg GST fusion protein + primary antibody dilution at 1) 1:5000 2) 1:10000