

Anti-PGP9.5 antibody [SM5228] (STJA0005228)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/IHC/IF
Host / Source	Mouse
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Monoclonal
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB (H, M, R) 1:1000-1:2000 IHC/IF (H, M, R) 1:1000-1:2000
Formulation	PBS with 0.02% sodium azide, 100 Mu g/ml BSA and 50% glycerol.
Isotype	IgG1
Storage Instruction	Store at -20C for up to one year, and avoid repeated freeze-thaw cycles.

TARGET INFORMATION


Gene ID	22223
Gene Symbol	Uchl1
UniProt ID	Uchl1_MOUSE

ADDITIONAL INFORMATION


Note STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.




Western blot analysis of PGP9.5. Sample: Protein treated by RIPA Lysis Buffer. Blocking buffer: 3% Nonfat dry milk in TBST, RT, 1h. Primary antibody: 1:1000, 4C overnight. Secondary antibody: HRP Goat Anti-mouse IgG, 1:3000, RT, 1h.




Immunohistochemistry analysis of PGP9.5. Sample: Human esophageal cancer, 4% PFA 12-24h. Antigen retrieval: Citrate buffer, 98C, 20 min. Blocking buffer: 3% BSA in PBS, RT, 30min. Primary antibody: 1:1000, 4C overnight. Secondary antibody: HRP Goat Anti-mouse IgG, 1:200 RT 1h.




Immunohistochemistry analysis of PGP9.5. Sample: Human stomach, 4% PFA 12-24h. Antigen retrieval: Citrate buffer, 98C, 20 min. Blocking buffer: 3% BSA in PBS, RT, 30min. Primary antibody: 1:1000, 4C overnight. Secondary antibody: HRP Goat Anti-mouse IgG, 1:200 RT 1h.




Immunohistochemistry analysis of PGP9.5. Sample: mouse brain, 4% PFA 12-24h. Antigen retrieval: Citrate buffer, 98C, 20 min. Blocking buffer: 3% BSA in PBS, RT, 30min. Primary antibody: 1:1000, 4C overnight. Secondary antibody: HRP Goat Anti-mouse IgG, 1:200 RT 1h.




Immunohistochemistry analysis of PGP9.5. Sample: mouse colon, 4% PFA 12-24h. Antigen retrieval: Citrate buffer, 98C, 20 min. Blocking buffer: 3% BSA in PBS, RT, 30min. Primary antibody: 1:1000, 4C overnight. Secondary antibody: HRP Goat Anti-mouse IgG, 1:200 RT 1h.



Immunohistochemistry analysis of PGP9.5. Sample: Rat brain, 4% PFA 12-24h. Antigen retrieval: Citrate buffer, 98C, 20 min. Blocking buffer: 3% BSA in PBS, RT, 30min. Primary antibody: 1:1000, 4C overnight. Secondary antibody: HRP Goat Anti-mouse IgG, 1:200 RT 1h.



Immunohistochemistry analysis of PGP9.5. Sample: Rat colon, 4% PFA 12-24h. Antigen retrieval: Citrate buffer, 98C, 20 min. Blocking buffer: 3% BSA in PBS, RT, 30min. Primary antibody: 1:1000, 4C overnight. Secondary antibody: HRP Goat Anti-mouse IgG, 1:200 RT 1h.



Immunohistochemistry analysis of PGP9.5. Sample: Rat stomach, 4% PFA 12-24h. Antigen retrieval: Citrate buffer, 98C, 20 min. Blocking buffer: 3% BSA in PBS, RT, 30min. Primary antibody: 1:1000, 4C overnight. Secondary antibody: HRP Goat Anti-mouse IgG, 1:200 RT 1h.